

# **Towards the Prediction of Decadal to Multi-Century Processes in a High-Throughput Climate System Model**

PI: Zhengyu Liu  
Center for Climatic Research  
University of Wisconsin-Madison

CoPIs:  
J. Kutzbach, UW-Madison  
R. Jacob, Argonne National Lab.  
C. Prentice, Max Planck Institute of Biogeochemistry  
J. Anderson, UW-Madison (formerly)

## **Climate Change, Climate Variability and Land-Vegetation Feedback: A Preliminary Report on FOAM Modeling**

The Fast Ocean-Atmosphere Model (FOAM) has been used to study the climate system. Three preliminary studies are carried out:

### 1) Holocene climate and climate variability:

The model is first tested against paleoclimate data. FOAM simulations of early to mid-Holocene climate (11,000 – 6,000 years ago) agree with most paleo observations of major monsoons (Liu et al., 2001; Harrison et al., 2001). The model also correctly simulates the change of ENSO variability through the Holocene. Based on the model simulation, it is proposed that the enhanced NH monsoon may contribute to the suppression of ENSO in the mid-Holocene (Liu et al., 1999; 2000).

### 2) Mechanism of interdecadal climate variability:

The model is used to understand the mechanism of decadal climate variability. In the Pacific, model simulations suggest that both the extratropical and tropical Pacific coupled ocean-atmosphere system can generate interdecadal climate variability. In addition, tropical-extratropical oceanic teleconnection seems to enhance the tropical decadal variability (Liu et al., 2001). In the Atlantic, model simulations suggest that the decadal variability can be generated within the tropical Atlantic ocean-atmosphere system, while North Atlantic Oscillation also affects the magnitude of tropical Atlantic variability (Liu and Wu, 2000; Wu and Liu, 2001).

### 3) Coupled climate-land vegetation system:

A sensitivity study suggests that deforestation in the Indonesian Archipelago, amplified by ocean-atmosphere feedback, could result a significant SST change in the tropical Indian Ocean (Delire et al., 2001). Furthermore, a dynamic land-vegetation model (LPJ) is being coupled to FOAM (Gallimore and Prentice, 2001), which will later be used to study environmental impact and feedback of land-ecosystem on climate.

## References

Delire C., P. Behling, M. T. Coe, J.A. Foley, R. Jacob, J. Kutzbach, Z. Liu and S. Vavrus, 2001: Response of coupled ocean-atmosphere system to deforestation in the Indonesian Archipelago. *Geophys. Res. Lett.*, **28**, 2081-2084

Gallimore and Prentice, 2001: Coupling a dynamic vegetation model to a climate model. in preparation.

Harrison, S., J. E. Kutzbach, Z. Liu et al.; 2001: North America monsoon in the mid-Holocene. *Clim. Dyn.*, submitted.

Liu, Z., R. Jacobs, J. Kutzbach, S. Harrison and J. Anderson, 1999: Asian monsoon impact on El Nino in the early Holocene. *PAGES Newsletter*, Vol. 7, No. 2, 16-17

Liu, Z., J. Kutzbach and L. Wu, 2000: Modeling climatic shift of El Nino variability in the Holocene. *Geophys. Res. Lett.*, **27**, 2265-2268.

Liu Z. and L. Wu, 2000: Tropical Atlantic Oscillation in a coupled GCM. *Atmos. Sci. Lett.*, **1**, <http://www.idealibrary.com/links/toc/asle/0/0/0>

Liu, Z., L. Wu, R. Gallimore and R. Jacob, 2001: Dual origins of Pacific interdecadal climate variability: the tropical mode and the North Pacific mode. GRL, submitted.

Liu, Z., S. Harrison, J. Kutzbach et al: 2001: Global monsoons in the Holocene and oceanic feedback. *J. Clim.*, submitted.